

Increased Expression of Cannabinoid CB₁ Receptors in Achilles Tendinosis

Emmelie Björklund^{1*}, Sture Forsgren², Håkan Alfredson³, Christopher J. Fowler¹

1 Pharmacology Unit, Department of Pharmacology and Clinical Neuroscience, Umeå University, Umeå, Sweden, **2** Section for Anatomy, Department of Integrative Medical Biology, Umeå University, Umeå, Sweden, **3** Sports Medicine Unit, Department of Surgical and Perioperative Sciences, Umeå University, Umeå, Sweden

Abstract

Background: The endogenous cannabinoid system is involved in the control of pain. However, little is known as to the integrity of the cannabinoid system in human pain syndromes. Here we investigate the expression of the cannabinoid receptor 1 (CB₁) in human Achilles tendons from healthy volunteers and from patients with Achilles tendinosis.

Methodology: Cannabinoid CB₁ receptor immunoreactivity (CB₁IR) was evaluated in formalin-fixed biopsies from individuals suffering from painful Achilles tendinosis in comparison with healthy human Achilles tendons.

Principal Findings: CB₁IR was seen as a granular pattern in the tenocytes. CB₁IR was also observed in the blood vessel wall and in the perineurium of the nerve. Quantification of the immunoreactivity in tenocytes showed an increase of CB₁ receptor expression in tendinosis tissue compared to control tissue.

Conclusion: Expression of cannabinoid receptor 1 is increased in human Achilles tendinosis suggesting that the cannabinoid system may be dysregulated in this disorder.

Citation: Björklund E, Forsgren S, Alfredson H, Fowler CJ (2011) Increased Expression of Cannabinoid CB₁ Receptors in Achilles Tendinosis. PLoS ONE 6(9): e24731. doi:10.1371/journal.pone.0024731

Editor: Naj Sharif, Alcon Research, Ltd., United States of America

Received: May 30, 2011; **Accepted:** August 16, 2011; **Published:** September 8, 2011

Copyright: © 2011 Björklund et al. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

Funding: The authors thank the Swedish Science Council (Grant no. 12158, medicine, to C.J. Fowler); the Research Funds of the Medical Faculty, Umeå University (to C.J. Fowler); the Faculty of Medicine at Umeå University, the Swedish National Centre for Research in Sports and the J.C. Kempe and Seth M. Kempe Memorial Foundations, Örnsköldsvik for financial support. The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.

Competing Interests: The authors have declared that no competing interests exist.

* E-mail: emmelie.bjorklund@pharm.umu.se

Introduction

It has been known for many years that cannabinoids are effective for the relief of a variety of types of pain [1], and activation of cannabinoid (CB) receptors by compounds such as nabilone have been shown to have clinical utility in a variety of pain states, including cancer pain, neuropathic pain, and fibromyalgia [2–4]. Data from studies on experimental animals indicate that the antinociceptive effects of cannabinoids are not only centrally mediated, but that spinal and peripheral CB receptors are involved [5].

The antinociceptive effects of exogenous CB receptor agonists raise the possibility that the endocannabinoid system is dysfunctional in pain states. There is evidence of this in animal models of pain [6–10], but little is known about the situation in human pain. To our knowledge, the only data so far reported are findings of an increased plasma concentration of the endocannabinoid anandamide in patients with complex regional pain syndrome compared to age- and sex-matched controls [11], a negative correlation between CB₁ receptor expression in pancreatic nerves and strong pain symptoms in patients with pancreatic cancer [12], and a positive correlation of suburothelial CB₁-immunoreactive nerve fibers with the pain score in painful bladder syndrome [13]. There is thus a need for more data on the status of the endocannabinoid system in human pain states.

The Achilles tendon is the strongest tendon in the body and the understanding of pain in this tendon is of utmost importance. When

there is chronic pain and impaired function in the Achilles tendon, the condition is referred to as tendinopathy. Clinically, the tendon thickens and becomes tender. Biopsies show changes in the appearance of the tenocytes, hypercellularity and neovascularization and eventually degenerative features [14–19]. When there are structural changes, the condition is commonly called tendinosis [20].

Achilles tendinosis most often occurs in the mid-portion of the tendon. Imaging with ultrasonography or magnetic resonance imaging (MRI) can be used to verify tendon abnormalities in the painful area [18,21]. The pain in chronic Achilles tendinosis remains an enigma. However, injections and mini-surgery to areas of neovascularization outside the tendon have been shown to reduce the pain seen in chronic Achilles tendinosis [22,23]. Although several molecular candidates have been identified and proposed as mediators of the pain in Achilles tendinosis [24], information on the cannabinoid system is lacking. In consequence, we have investigated the expression of CB₁ receptors in human Achilles tendons and whether there is a change in expression in Achilles tendinosis.

Methods

Ethics Statement

The Committee of Ethics at the Faculty of Medicine, Umeå University and the Regional Ethical Review Board in Umeå approved the project (04-157 M). Participants gave verbal consent

to participate in the research after reading an explanatory statement and receiving a verbal summary of the project. The Review Board approved the verbal consent procedure. When approval from patients had been obtained, specimens were collected and taken care of for the research and documentation was made. This form of approval and documentation is in accordance with paragraph 16 and 17 of the Ethical review act in Sweden. All procedures followed the principles of the Declaration of Helsinki.

Individuals

This investigation included tissue samples from 24 individuals: 11 males and 13 females (mean age: 48 years, range 21–70 years; male mean age: 44 years, range 28–70 years; female mean age: 52 years, range 21–68 years). The samples were from a group of patients suffering from Achilles tendinosis and from healthy controls. The Achilles tendinosis group ($n = 17$) were suffering from chronic painful mid-portion Achilles tendinosis. This group consisted of 8 males and 9 females (mean age: 51, range 28–70 years; male mean age: 45 years, range 28–70; female mean age: 57 years, range 47–68 years). Tendinosis was verified by ultrasonography. The control group consisted of individuals ($n = 7$) with no history of pain symptoms from their Achilles tendons (mean age: 41 years, range 21–47 years; 3 males, mean age 41 years, range 39–46 years; 4 females, mean age: 41, range 21–47 years). Ultrasonography showed normal tendons. All patients were otherwise healthy and non-smokers.

Sampling, fixation and sectioning

In the tendinosis group, biopsies were taken during surgical treatment. Under local anaesthesia (4–5 ml pilocaine hydrochloride, 10 mg/ml; AstraZeneca, Södertälje, Sweden), tendon tissue (macroscopically abnormal) was taken from the ventral part of the Achilles tendon through a longitudinal incision lateral to the tendon mid-portion. The biopsies contained tendon tissue proper, and to varying degrees, outer parts of the tendon (paratendinous connective tissue). The tissue samples were approximately 2 mm wide and 1–5 mm long and were taken from different depths of the tendon. From the control group, samples of the same size were collected from the dorsal part of the tendon using a longitudinal plain incision under local anaesthesia. The dorsal part of the tendon was chosen for ethical and practical reasons.

Directly after the surgical procedures, the samples were fixed overnight at 4°C in a solution of 4% formaldehyde in 0.1 M phosphate buffer, pH 7.0, followed by thorough washing in Tyrode's solution containing 10% sucrose, at 4°C overnight. The samples were then mounted on thin cardboard in OCT embedding medium (Miles Laboratories, Naperville, IL, USA) and frozen at –80°C until sectioning.

For the immunofluorescence and morphologic investigations, series of 7 µm thick sections from both groups of samples were cut using a cryostat (Leica Microsystem CM3050S, Heidelberg, Germany). The sections were mounted on slides, pre-coated with crome-alun gelatine, and the sections were then dried and thereafter processed for immunohistochemistry. Staining with hematoxylin-eosin was carried out to delineate tissue morphology. Reference tissues (human colonic and rat dorsal root ganglion tissue), available at the laboratory, were examined in parallel. They had been fixed and further processed in the same way as the tendon samples.

Immunofluorescence

Sections were initially treated with acid potassium permanganate for 2 min to enhance the visualization of specific immuno-

fluorescence reaction sites [25]. The sections were then rinsed in 0.01 M phosphate-buffered saline (PBS) pH 7.4 containing 0.1% sodium azide as preservative, for 3×5 min. After these procedures, incubation for 20 min in a 1% solution of detergent Triton X-100 (Kebo Lab, Stockholm, Sweden) in 0.01 M PBS, and rinsing for 3×5 min in PBS were performed. The sections were then incubated in 5% normal swine serum in PBS supplemented with 0.1% bovine serum albumin (BSA) for 15 min in a humid environment in room temperature. Thereafter followed incubation with the primary antibody, diluted in PBS with BSA. The incubation was performed in a humid environment and proceeded for 60 min at 37°C. The sections were then washed in PBS for 3×5 min prior to another incubation in normal swine serum for 15 min in a humid environment in room temperature, followed by incubation with the secondary antibody. This antibody corresponded to tetramethylrhodamine isothiocyanate (TRITC)-conjugated swine antirabbit IgG (DakoCytomation, Glostrup, Denmark). The secondary antibody was diluted in PBS with BSA (1:40). Incubation proceeded for 30 min at 37°C in a humid environment. A Zeiss Axioskop II microscope equipped with an Olympus DP70 digital camera was used for examination of the sections.

For semiquantitative determination of the intensity of CB₁ receptor immunoreactivity in the control and Achilles tendinosis tendons, two of the researchers (EB, SF), who were blinded to the clinical data for the patients, independently scored the CB₁ expression in the sections under the fluorescence microscope. The tenocytes were scored for immunoreactive intensity (0–3 where 0 is absent and 3 is high), and the average value was taken. This method of scoring has previously been used in other published studies by co-author Forsgren [26,27], including studies using the immunofluorescence technique on tendon tissue [28]. For the CB₁ receptor immunoreactivity scores returned by the two investigators, an intraclass coefficient analysis using a two-way mixed effects model for the 24 scores gave a Cronbach's alpha reliability coefficient of 0.79, suggesting that the inter-rater reliability is acceptable.

Antibodies and control stainings

The antibodies used were against CB₁ and PGP9.5. Both are rabbit antibodies. The CB₁ antibody (rabbit polyclonal; ab23703; Abcam, Cambridge, UK) had been raised against the C-terminal amino acid 461–472 of human CB₁. Concentrations used were 1:20–1:100. For the experiments comparing immunoreactivity for controls and tendinosis patients, a dilution of 1:50 was used. The antibody to PGP9.5 (rabbit polyclonal; code 7863-0504; Biogenesis, Poole, UK), was used at a dilution of 1:100, and had been raised against native brain PGP9.5.

For control purposes, sections were processed with PBS/BSA instead of primary antibodies. To confirm further the specificity, the primary CB₁ antibody was pre-absorbed overnight at 4°C with its immunogenic peptide (20–100 µg/ml; ab50542; Abcam, Cambridge, UK) prior to incubation on the sections. The PGP9.5 antibody has been used in numerous studies in the laboratory [29,30] and in numerous studies by others for the purpose of demarcating nerve fibres.

Statistics

The two-tailed Mann-Whitney test was undertaken using the Statistical package built into the GraphPad Prism 5 computer programme for the Macintosh (GraphPad Software Inc., San Diego, CA, USA). The intraclass coefficient analysis was conducted using IBM SPSS Statistics 19 for the Macintosh (IBM Inc., Somers, NY, USA).

Results

CB₁ receptor immunoreactivity (CB₁IR) in the normal Achilles tendon and in reference tissue

Figure 1 shows the CB₁IR for normal Achilles tendons, in which the tendon cells (tenocytes) show the characteristic elongated appearance [18,19,31] and, for comparative purposes, that for human colon and the rat dorsal root ganglion. For all three cases, CB₁IR was seen. For the tendons, CB₁IR was observed for the tenocytes (Fig. 1A) and in blood vessel walls (not shown). The reactions in the tenocytes showed a punctuate appearance along the length of the cells (Fig. 1A). Preabsorption with the corresponding peptide eliminated the immunostainings (Fig. 1B), as did control stainings when the primary antibody was omitted (data not shown). As positive controls, CB₁IR was investigated in the human colon and the dorsal root ganglia. For the human colon, distinct specific reactions were seen for cells of the mucosa and submucosa and in cells in the epithelial layer (Fig. 1C), in agreement with the literature [32]. Distinct specific reactions were also seen for neuronal perikarya of the dorsal root ganglion. The reactions occurred as granular intracellular reactions (Fig. 1E). The presence of intracellular CB₁IR can also be seen in the literature [33]. For both human colon and the rat dorsal root ganglia, preabsorption with the corresponding peptide eliminated the immunostainings (Fig. 1D and F).

CB₁IR in Achilles tendinosis

The CB₁IR pattern in tenocytes of samples obtained from patients with Achilles tendinosis is shown in low magnification in (Fig. 2A–B), where Fig. 2B shows that preincubation with the antigen prevents all the immunoreactive staining. At a higher magnification, a granular pattern of immunoreactivity was seen for the tenocytes. That was especially the situation for rounded/swollen tenocytes and tenocytes with a wavy appearance. These abnormally-formed tenocytes are a characteristic of Achilles tendinosis [16,19]. The tenocyte reactions appeared overall to be the strongest in the tendinosis specimens.

CB₁IR in blood vessel walls and nerve structures

CB₁IR was also observed in blood vessel walls and in the perineurium of nerves in the samples, especially in the tendinosis samples. For the blood vessel walls (confirmed by htx-eosin staining Fig. 3A), CB₁IR was in the form of fine pointed reactions (Fig. 3B). They were seen for small vessels but not for large vessels. Preabsorption with the immunising peptide confirmed specificity of the antibody (data not shown).

Nerve fascicles were demarcated via showing a distinct PGP9.5 immunoreaction (Fig. 3C). CB₁IR was detected in the perineurium of the nerve fascicles. No such reactions were, on the other hand, detected within the interior of the nerve fascicles (Fig. 3D). The nerve related immunoreactions were abolished following preabsorption with the antigen peptide (data not shown).

Semiquantitative comparison of tenocyte CB₁IR in controls and patients with Achilles tendinosis

A semiquantitative analysis of the reaction intensities for the tenocytes was performed. CB₁ was scored in tenocytes from a total of 24 patients either suffering from chronic Achilles tendinosis or having clinically pain-free Achilles tendons. Difference in CB₁ expression between groups was statistically significant ($p < 0.05$, Mann-Whitney U test) (Fig. 4). It should be noted that the control patient with the highest CB₁IR was presumably asymptomatic at the time of biopsy, but that the tendon showed pathology




Figure 1. Immunofluorescence for CB₁ in normal Achilles tendon and in reference tissue. Panels show sections processed for CB₁ (a,c,e) and with CB₁ antibody preabsorbed with CB₁ antigen (b,d,f). Immunoreactions are shown in the elongated tenocytes in (a) but not in (b). Arrows indicate tenocytes. Immunoreactions are also seen in cells of the mucosa and submucosa (arrows) and the epithelial layer (arrowheads) of human colon in (c) but not in (d), and in cell bodies of a rat dorsal root ganglion in (e) but not in (f) (arrows). Asterisks in similar region in (c) and (d). Original magnification $\times 40$. doi:10.1371/journal.pone.0024731.g001

consistent with Achilles tendinopathy, thus questioning whether this sample is a true control. Exclusion of this sample from the data set increased the level of significance to $p < 0.005$.

Discussion

In this study we demonstrate, for the first time, that CB₁IR is detectable within the tenocytes of the human Achilles tendon. Immunoreactions were also seen in the walls of small blood vessel walls and in the perineurium of nerves. On the other hand, whilst CB₁IR was detectable in the neuronal perikarya of the rat spinal ganglion, CB₁IR was not detectable in the nerves of tendon tissue.

The antibody used for these studies, raised against the C-terminal amino acid 461–472 sequence of human CB₁, has been found previously to produce the appropriate staining in brain samples, but not to stain samples from CB₁ knockout mice [34,35]. The antibody has been used, by us and others, to investigate CB₁ receptor expression pattern in human bladder [36] and fallopian tube [37], and in colorectal and prostate cancer biopsies [34,35]. We noted that immunoreactivity was present for non-neuronal cells and that it was not limited to the cell plasma membranes, but that intracellular reactions were also seen. Similarly we noted in the present study that the CB₁IR was detectable intracellularly in the neuronal cell perikarya and in the cells of the mucosa/submucosa and the epithelial cells of the colon. Although at first sight this may seem strange for a G-protein coupled receptor, it is known that the majority of the endogenous CB₁ receptors do not

Achilles tendon tissue as such, the existing innervations being present in the outer part of the tendon (the peritendinous tissue) and to a very small extent in the connective tissue septa [30]. These findings may suggest that biochemical mediators, such as nerve signal substances, produced in the tendon tissue, can be involved in the aetiology and pathogenesis of chronic tendon pain in tendinosis [47]. A “biochemical model” of this type, involving other locally produced signal substances, has also previously been considered [48]. This means that the pain is not only structural in origin. To what extent such a model is related to what we see here, i.e. an up-regulation of CB₁IR for tenocytes in tendinosis tendons, the CB₁ immunoreactions being especially obvious for tenocytes with an abnormal morphology, remains to be further explained. In any case, our observations show that the cannabinoid receptor system in the Achilles tendon is mainly related to the tendon cells and not being confined to the nerve fibres.

In the present study, the robust and reliable semiquantitative determination method used, also found that there was a higher CB₁ receptor expression level in the tenocytes of patients with Achilles tendinosis than in the control group, suggesting that the cannabinoid system may be dysregulated in this disorder. It is possible that the increased expression may be an adaptive consequence to a loss of local endocannabinoid signalling, and it would clearly be of interest to investigate local endocannabinoid levels in microdialysis studies from this patient group.

An alternative explanation is that the increased CB₁ receptor expression is due to changes in its cellular regulation produced by the local environment in the disorder. There is evidence in the literature that inflammatory processes can affect expression of CB₁ receptors in some cases, but not others: Izzo et al. [49] for example demonstrated that croton oil-induced intestinal inflammation resulted in a three-fold increase in CB₁ receptor expression in the jejunum. In contrast, cystitis produced by intravesical administration of acrolein did not affect bladder CB₁ investigation

at the time points chosen [50]. CB₁ and CB₂ receptors can be regulated by both pro- and anti-inflammatory cytokines [51], and a good example of this is the ability of interleukin-4 (IL-4) to increase CB₁ receptor expression in lymphocytes by a Stat-6 mediated pathway [52,53]. Tenocytes in primary culture express interleukin-4 (IL-4) receptors [45]. However, inflammation presumably occurs only at the beginning of tendinosis, and there are no signs of inflammation at the time when the biopsy samples used here were taken, so that if the increased expression found in the present study emanates from the early inflammation, it would have to be a very long-lasting effect, or due to the residual presence of a factor long after resolution of the inflammation. This does not, of course, rule out the possibility that other factors, i.e. not associated with inflammation, are involved in the regulation of CB₁ receptor expression in the tendons. Experiments in both patient tissue and animal models of Achilles tendinopathy [54] are clearly needed to explore further the mechanisms behind the increased CB₁ receptor expression in Achilles tendinosis.

In conclusion, the present study has demonstrated the expression of CB₁ receptors in Achilles tenocytes, thereby adding to the growing list of non-neuronal cell types that express these receptors, and that their expression is increased in tendinosis. These data underline the need for further studies on endocannabinoid signalling in human pain conditions.

Acknowledgments

The authors are grateful to Ulla Hedlund for expert technical assistance.

Author Contributions

Conceived and designed the experiments: EB SF CJF. Performed the experiments: EB. Analyzed the data: EB SF. Contributed reagents/materials/analysis tools: HA. Wrote the paper: EB SF CJF.

References

- Reynolds J (1890) On the therapeutical uses and toxic effects of *cannabis indica*. *Lancet* 135: 637–638.
- Frank B, Serpell M, Hughes J, Matthews J, Kapur D (2008) Comparison of analgesic effects and patient tolerability of nabilone and dihydrocodeine for chronic neuropathic pain: randomised, crossover, double blind study. *BMJ* 336: 199–201.
- Maida V, Ennis, Irani S, Corbo M, Dolzhykov M (2008) Adjunctive nabilone in cancer pain and symptom management: a prospective observational study using propensity scoring. *J Support Oncol* 6: 119–124.
- Skrabek RQ, Galimova L, Ethans K, Perry D (2008) Nabilone for the treatment of pain in fibromyalgia. *J Pain* 9: 164–173.
- Guindon J, Hohmann AG (2009) The endocannabinoid system and pain. *CNS Neurol Disord Drug Targets* 8: 403–421.
- Mitirattanakul S, Ramakul N, Guerrero A, Matsuka Y, Ono Y, et al. (2006) Site-specific increases in peripheral cannabinoid receptors and their endogenous ligands in a model of neuropathic pain. *Pain* 126: 102–114.
- Khasabova I, Khasabov S, Harding-Rose C, Coicou L, Seybold B, et al. (2008) A decrease in anandamide signaling contributes to the maintenance of cutaneous mechanical hyperalgesia in a model of bone cancer pain. *J Neurosci* 28: 11141–11152.
- Hong S, Fan J, Kemmerer E, Evans S, Li Y, et al. (2009) Reciprocal changes in vanilloid (TRPV1) and endocannabinoid (CB1) receptors contribute to visceral hyperalgesia in the water avoidance stressed rat. *Gut* 58: 202–210.
- Hoot MR, Sim-Selley LJ, Poklis JL, Abdullah RA, Scoggins KL, et al. (2010) Chronic constriction injury reduces cannabinoid receptor 1 activity in the rostral anterior cingulate cortex of mice. *Brain Res* 1339: 18–25.
- Bishay P, Schmidt H, Marian C, Häussler A, Wijnvoord N, et al. (2010) R-flurbiprofen reduces neuropathic pain in rodents by restoring endogenous cannabinoids. *PLoS ONE* 5: e10628.
- Kaufmann I, Hauer D, Hüge V, Vogeser M, Campolongo P, et al. (2009) Enhanced anandamide plasma levels in patients with complex regional pain syndrome following traumatic injury: a preliminary report. *Eur Surg Res* 43: 325–329.
- Michalski C, Oti F, Erkan M, Sauliunaite D, Bergmann F, et al. (2008) Cannabinoids in pancreatic cancer: correlation with survival and pain. *Int J Cancer* 122: 742–750.
- Mukerji G, Yiangou Y, Agarwal SK, Anand P (2010) Increased cannabinoid receptor 1-immunoreactive nerve fibers in overactive and painful bladder disorders and their correlation with symptoms. *Urology* 75: 1514 e1515–1520.
- Äström M, Rausing A (1995) Chronic Achilles tendinopathy. A survey of surgical and histopathologic findings. *Clin Orthop Relat Res* 316: 151–164.
- Movin T, Gad A, Reinholt FP, Rolf C (1997) Tendon pathology in long-standing achillobodynia. Biopsy findings in 40 patients. *Acta Orthop Scand* 68: 170–175.
- Khan KM, Cook JL, Bonar F, Harcourt P, Åström M (1999) Histopathology of common tendinopathies. Update and implications for clinical management. *Sports Med* 27: 393–408.
- Maffulli N, Testa V, Capasso G, Ewen SW, Sullo A, et al. (2004) Similar histopathological picture in males with Achilles and patellar tendinopathy. *Sports Sci Sports Exerc* 36: 1470–1475.
- Shalabi A, Movin T, Krisoffersen-Wiberg M, Aspelin P, Svensson L (2005) Reliability in the assessment of tendon volume and intratendinous signal of the Achilles tendon on MRI: a methodological description. *Knee Surg Sports Traumatol Arthrosc* 13: 492–498.
- Bjur D, Danielson P, Alfredson H, Forsgren S (2008) Immunohistochemical and in situ hybridization observations favor a local catecholamine production in the human Achilles tendon. *Histol Histopathol* 23: 197–208.
- Alfredson H (2005) The chronic painful Achilles and patellar tendon: research on basic biology and treatment. *Scand J Med Sports* 15: 252–259.
- Alfredson H, Öhberg L, Forsgren S (2003) Is vasculo-neural ingrowth the cause of pain in chronic Achilles tendinosis? An investigation using ultrasonography and colour Doppler, immunohistochemistry, and diagnostic injections. *Knee Surg Sports Traumatol Arthrosc* 11: 334–338.
- Alfredson H (2011) Ultrasound and Doppler-guided mini-surgery to treat midportion Achille tendinosis: results of a large material and a randomised study comparing two scraping techniques. *Br J Sports Med* 45: 407–410.
- Alfredson H, Öhberg L (2005) Sclerosing injections to areas of neovascularisation reduce pain in chronic Achilles tendinopathy: A double-blind randomized controlled trial. *Knee Surg Sports Traumatol Arthrosc* 13: 338–344.
- Riley G (2008) Tendinopathy- from basic science to treatment. *Nature clinical practice* 4: 82–89.

25. Hansson M, Forsgren S (1995) Immunoreactive atrial and brain natriuretic peptides are co-localized in Purkinje fibres but not in the innervation of the bovine heart conduction system. *Histochem J* 27: 222–230.
26. Grimsholm O, Guo Y, Ny T, Forsgren S (2008) Expression patterns of neurotrophins and neurotrophin receptors in articular chondrocytes and inflammatory infiltrates in knee joint arthritis. *Cells Tissues Organs* 188: 299–309.
27. Johansson M, Norrgård O, Forsgren S (2007) Study of expression patterns and levels of neurotrophins and neurotrophin receptors in ulcerative colitis. *Inflamm Bowel Dis* 13: 398–409.
28. Bjur D, Alfredson H, Forsgren S (2009) Presence of the neuropeptide Y1 receptor in tenocytes and blood vessel walls in the human Achilles tendon. *Br J Sports Med* 43: 1136–1142.
29. Andersson G, Danielson P, Alfredson H, Forsgren S (2007) Nerve-related characteristics of ventral paratendinous tissue in chronic Achilles tendinosis. *Knee Surg Sports Traumatol Arthrosc* 15: 1272–1279.
30. Bjur D, Alfredson H, Forsgren S (2005) The innervation pattern of the human Achilles tendon: studies of the normal and tendinosis tendon with markers for general and sensory innervation. *Cell Tissue Res* 320: 201–206.
31. Khan KM, Cook JL, Bonar F, Harcourt P, Astrom M (1999) Histopathology of common tendinopathies. Update and implications for clinical management. *Sports Med* 27: 393–408.
32. Marquéz L, Suárez J, Iglesias M, Bermudez-Silva FJ, Rodríguez de Fonseca F, et al. (2009) Ulcerative colitis induces changes on the expression of the endocannabinoid system in the human colonic tissue. *PLoS ONE* 4: e6893.
33. Bridges D, Rice ASC, Egertová M, Elphick MR, Winter J, et al. (2003) Localisation of cannabinoid receptor 1 in rat dorsal root ganglion using in situ hybridisation and immunohistochemistry. *Neuroscience* 119: 803–812.
34. Chung SC, Hammarsten P, Josefsson A, Stattin P, Granfors T, et al. (2009) A high cannabinoid CB₁ receptor immunoreactivity is associated with disease severity and outcome in prostate cancer. *Eur J Cancer* 45: 174–182.
35. Gustafsson SB, Palmqvist R, Henriksson ML, Dahlin AM, Edin S, et al. (2011) High tumour cannabinoid CB₁ receptor immunoreactivity negatively impacts disease-specific survival in stage II microsatellite stable colorectal cancer. *PLoS ONE*;in press.
36. Tyagi V, Phillips B, Su R, Smaldone MC, Erickson VL (2009) Differential expression of functional cannabinoid receptors in human bladder detrusor and urothelium. *J Urol* 181: 1932–1938.
37. Horne AW, Phillips JA, 3rd, Kane N, Lourenco PC, McDonald SE (2008) CB1 expression is attenuated in Fallopian tube and decidua of women with ectopic pregnancy. *PLoS ONE* 3: e3969.
38. Rozenfeld R, Devi LA (2008) Regulation of CB₁ cannabinoid receptor trafficking by the adaptor protein AP-3. *FASEB J* 22: 2311–2322.
39. McIntosh HH, Song C, Howlett AC (1998) CB₁ cannabinoid receptor: cellular regulation and distribution in N18TG2 neuroblastoma cells. *Brain Res Mol Brain Res* 53: 163–173.
40. Ong WY, Mackie K (1999) A light and electron microscopic study of the CB1 cannabinoid receptor in primate brain. *Neuroscience* 92: 1177–1191.
41. Scavone JL, Mackie K, Van Bockstaele EJ (2010) Characterization of cannabinoid-1 receptors in the locus coeruleus: relationship with mu-opioid receptors. *Brain Res* 1312: 18–31.
42. Brailoiu GC, Oprea TI, Zhao P, Abood ME, Brailoiu E (2011) Intracellular CB₁ cannabinoid receptors are activated by anandamide. *JBC*;in press.
43. Andersson G, Danielson P, Alfredson H, Forsgren S (2008) Presence of substance P and the neurokinin-1 receptor in tenocytes of the human Achilles tendon. *Regul Pept* 150: 81–7.
44. Bjur D, Danielson P, Alfredson H, Forsgren S (2008) Presence of a non-neuronal cholinergic system and occurrence of up- and down-regulation in expression of M2 muscarinic acetylcholine receptors: new aspects of importance regarding Achilles tendon tendinosis (tendinopathy). *Cell Tissue Res* 331: 385–400.
45. Courneya JP, Luzina IG, Zeller CB, Rasmussen JF, Bocharov A, et al. (2010) Interleukins 4 and 13 modulate gene expression and promote proliferation of primary human tenocytes. *Fibrogenesis Tissue Repair* 3: 9.
46. Scott A, Alfredson H, Forsgren S (2008) VGLuT2 expression in painful Achilles and patellar tendinosis: evidence of local glutamate release by tenocytes. *J Orthop Res* 26: 197–208.
47. Danielson P (2009) Reviving the “biochemical” hypothesis for tendinopathy: new findings suggests the involvement of locally produced signal substances. *Br J Sports Med* 43: 265–268.
48. Khan KM, Cook JL, Maffulli N, Kannus P (2000) Where is the pain coming from in tendinopathy? It may be biochemical, not only structural, in origin. *Br J Sports Med* 34: 81–83.
49. Izzo A, Fezza F, Capasso F, Bisogno T, Pinto L, et al. (2001) Cannabinoid CB₁-receptor mediated regulation of gastrointestinal motility in mice in a model of intestinal inflammation. *Br J Pharmacol* 134: 563–570.
50. Merriam FV, Wang Z-y, Guerios SD, Bjorling DE (2008) Cannabinoid receptor 2 is increased in acutely and chronically inflamed bladder of rats. *Neurosci Letts* 445: 130–134.
51. Jean-Gilles L, Gran B, Constantinescu CS (2010) Interaction between cytokines, cannabinoids and the nervous system. *Immunobiology* 215: 606–610.
52. Börner C, Bedini A, Höllt V, Kraus J (2008) Analysis of promoter regions regulating basal and interleukin-4-inducible expression of the human CB1 receptor gene in T lymphocytes. *Mol Pharmacol* 73: 1013–1019.
53. Börner C, Höllt V, Kraus J (2007) Activation of human T cells induces upregulation of cannabinoid receptor type 1 transcription. *Neuroimmunomodulation* 14: 281–286.
54. Andersson G, Forsgren S, Scott A, Gaida JE, Stjernfeldt JE, et al. (2011) Tenocyte hypercellularity and vascular proliferation in a rabbit model of tendinopathy: contralateral effects suggest the involvement of central neuronal mechanisms. *Br J Sports Med* 45: 399–406.